

A canonical curve of genus 17

Israel Moreno-Mejía

March 3, 2013

Abstract

We compute equations for a Hurwitz curve of genus 17 and we conclude that the canonical ideal of any Hurwitz curve of genus 14 or 17 is generated by quadrics.¹

1 Introduction

The curve alluded in the title is one of the 2 Hurwitz curves of genus 17 and the aim of this paper is to show that the ideal of the canonical embedding of this curve is generated by quadrics (see Theorem 10.1). A similar conclusion is reached in the genus 14 case (see Theorem 11.1). The proof is an application of *Petri's Analysis of the Canonical Ideal* (see for instance [9],[1],[8] or [10]) and makes use of an explicit basis for the space of quadrics in the ideal of the curve. To compute this basis we follow the method used in [7] for the case of the Hurwitz curves of genus 14; that is, we study the action of the curve's automorphism group G on the global sections of tensor powers of the canonical sheaf (see §5). Although calculations in the genus 17 case are similar to those in [7], some computations turned out to be very slow: namely, those of the matrices (34) and (40) and somehow the application of Petri's Analysis of the Canonical Ideal. One can convince oneself from the calculation of the quadrics that, it is possible to obtain equations for the remaining genus 17 and genus 14 Hurwitz curves by applying appropriate elements of the absolute Galois group $Gal(\bar{\mathbb{Q}}, \mathbb{Q})$ to the coefficients of the equations of the curves considered here. In the genus 14 case, this had already been observed in [12]. It follows that the canonical ideal of any genus 14 or

¹2000 MSC 14H37,14H45.

genus 17 Hurwitz curve is generated by quadrics. Most calculations were carried out with the help of Gap [4] and ParGap [3]. The matrices (34) and (40) can be obtained from me in case someone needs to compute explicitly this basis of quadrics and wants to save some time.

Very little is known about the genus 17 Hurwitz curves and the information available to us about their automorphism group comes from [2] and [11]. One could highlight the fact that there is only one Hurwitz group G of genus 17 and that it can be realized as the automorphism group of 2 non isomorphic genus 17 curves X_1 and X_2 (see Section 4 for the definition of X_i). Unlike the Hurwitz groups of genus 3, 7 and 14 our group G is not simple and has a normal subgroup N isomorphic to $(\mathbb{Z}/2\mathbb{Z})^3$. One has that G/N is $PSL(2, \mathbb{F}_7)$ and $X_1/N \cong X_2/N$ is the Hurwitz curve of genus 3.

We remark that it is not known a way to represent a general curve of genus g by means of polynomial equations, except for small values of g (see [5] Chap.6); and that although for some genera it is possible to give equations of special curves, it appears that for the genus 14 and genus 17 cases the Hurwitz curves are the only ones so far for which equations defining them can be computed.

2 Notation

We keep our notation as in [7]. Given a curve X and $h \in \text{Aut}(X)$ with fixed point set $\text{fix}(h) = \{p_1, \dots, p_u\}$ then

$$d_X h = (dh_{p_1}, \dots, dh_{p_u}),$$

where dh_{p_i} is the eigenvalue of h acting on K_{X,p_i} , where K_{X,p_i} is the fibre of the canonical line bundle of X at p_i .

If G is a finite group then $W_1, \dots, W_s$ denote its complex irreducible representations and $\chi_1, \dots, \chi_s$ denote the corresponding characters. Given a $\mathbb{C}G$ -module M we write M_{W_i} for the sum of all $\mathbb{C}G$ -submodules of M isomorphic to W_i . The projection $M \rightarrow M_{W_i}$ is denoted by π_{W_i} . For $h \in G$, $\rho(h)_{\nu_j}$ denotes the projection from M to the ν_j -eigenspace of h in M . Throught this paper we denote by G to the Hurwitz group of genus 17.

3 The automorphism group

In terms of generators and relations our group G has the following two inequivalent presentations (see [11])

$$G = \langle P, Q \mid P^7 = Q^8 = (QP^2)^3 = (QP^3)^2 = Q^4PQ^4P^4Q^4P^2 = 1 \rangle \quad (1)$$

$$\cong \langle P, Q \mid P^7 = Q^8 = (QP^2)^3 = (QP^3)^2 = Q^4PQ^4P^2Q^4P^4 = 1 \rangle. \quad (2)$$

Each presentation induces a pair of generators U, T given by

$$U = P^2Q, \quad (3)$$

$$T = P^3Q, \quad (4)$$

such that $U^2 = T^3 = (T^2U)^7 = 1$ from which one deduce that G is a Hurwitz group. Moreover, using an isomorphism between the two presentations one has two generating triplets (U, T, T^2U) , (U', T', T'^2U') such that T^2U and T'^2U' belong to different conjugacy classes. The outer automorphism group of G has order 2 and the two conjugacy classes of elements of order 7 are fixed under the action of the outer automorphisms of G . This gives rise to non isomorphic genus 17 curves X_1 and X_2 with automorphism group G (see for instance Theorem 2.17 in [13]).

In this paper we have chosen the presentation (1) as our definition of G . Using Gap one can check that the following gives us a representation of G as a permutation group

$$P = (1, 13, 2, 11, 4, 5, 8)(3, 10, 6, 14, 7, 9, 12), \quad (5)$$

$$Q = (1, 7, 3, 4)(2, 11, 13, 9, 6, 14, 10, 5), \quad (6)$$

that we use to represent some elements of G . Since Gap some times produces random character tables for a given group, we fix here the character table of G (see Table 1) that we used in our calculations. Representatives of the conjugacy classes are shown in Table 2 together with the size of the normalizer group in G of the cyclic group generated by a representative.

Table 1: Character Table of G

	1A	2A	4A	2B	4B	3A	6A	8A	8B	7A	7B
χ_1	1	1	1	1	1	1	1	1	1	1	1
χ_2	3	3	-1	-1	-1	0	0	1	1	β_3	β_1
χ_3	3	3	-1	-1	-1	0	0	1	1	β_1	β_3
χ_4	6	6	2	2	2	0	0	0	0	-1	-1
χ_5	7	7	-1	-1	-1	1	1	-1	-1	0	0
χ_6	7	-1	3	-1	-1	1	-1	1	-1	0	0
χ_7	7	-1	-1	-1	3	1	-1	-1	1	0	0
χ_8	8	8	0	0	0	-1	-1	0	0	1	1
χ_9	14	-2	2	-2	2	-1	1	0	0	0	0
χ_{10}	21	-3	1	1	-3	0	0	-1	1	0	0
χ_{11}	21	-3	-3	1	1	0	0	1	-1	0	0

Here $\beta_j = \zeta^j + \zeta^{2j} + \zeta^{4j}$, $\zeta = e^{2\pi i/7}$.

Table 2: Representatives of the conjugacy classes of G

Conjugacy class	Representative	$N_G(< h >)$
1A	Identity	1344
2A	(4, 7)(5, 9)(8, 12)(10, 13)	192
4A	(4, 5, 7, 9)(8, 10, 12, 13)	64
2B	(2, 6)(4, 9)(5, 7)(8, 13)(10, 12)(11, 14)	16
4B	(1, 3)(4, 5, 7, 9)(8, 13, 12, 10)(11, 14)	64
3A	(2, 4, 5)(6, 7, 9)(8, 10, 11)(12, 13, 14)	12
6A	(1, 3)(2, 4, 5)(6, 7, 9)(8, 13, 11, 12, 10, 14)	12
8A	(2, 4, 14, 12, 6, 7, 11, 8)(5, 13, 9, 10)	32
8B	(1, 3)(2, 4, 11, 12, 6, 7, 14, 8)(5, 10)(9, 13)	32
7A	(1, 2, 4, 8, 13, 11, 5)(3, 6, 7, 12, 10, 14, 9)	21
7B	(1, 8, 5, 4, 11, 2, 13)(3, 12, 9, 7, 14, 6, 10)	21

Table 3: The characteristic polynomials of $h|_{W_i}$

$h \in 7B$	
W_2	$(\lambda - \zeta)(\lambda - \zeta^2)(\lambda - \zeta^4)$
W_3	$(\lambda - \zeta^3)(\lambda - \zeta^5)(\lambda - \zeta^6)$
W_4	$(\lambda^7 - 1)/(\lambda - 1)$
W_5, W_6, W_7	$(\lambda^7 - 1)$
W_8	$(\lambda^7 - 1)(\lambda - 1)$
W_9	$(\lambda^7 - 1)^2$
W_{10}, W_{11}	$(\lambda^7 - 1)^3$

Here $\zeta = e^{2i\pi/7}$.

4 The action on the canonical line bundle

One can characterize the Hurwitz curves X_1 and X_2 in terms of the action of an element of order 7 on their canonical line bundles K_{X_j} , $j=1,2$. Let $d_{X_j}h$ be as defined in §2.

Lemma 4.1. *Let $h \in 7B$. For $j = 1, 2$ we can assume that $d_{X_j}h = (\zeta^{3^j-1}, \zeta^{3^j-1 \cdot 2}, \zeta^{3^j-1 \cdot 4})$, where $\zeta = e^{2i\pi/7}$.*

The proof is similar to that of Lemma 2.2 in [7].

So, from now on we denote by X_1 to the curve such that the action of a representative h_{7B} of the conjugacy class 7B gives the vector $d_{X_1}h_{7B} = (\zeta^1, \zeta^2, \zeta^4)$ and by X_2 to the curve such that $d_{X_2}h_{7B} = (\zeta^3, \zeta^5, \zeta^6)$. Now one can use [6, Theorem 1] to complete the Table 4, and then compute the multiplicities $a_i(d)$ of the representations W_i , $i = 1, \dots, 11$, in $H^0(X_j, K_{X_j}^d)$ (see Tables 5 and 6) by using the Atiyah–Bott fixed-point theorem and the character table of G .

Table 4: The fixed points and $d_{X_j}(h)$

Conjugacy class of h	Number of fixed points in X_j	$d_{X_j}(h)$
2A	0	—
4A	0	—
2B	8	$(-1, -1, -1, -1, -1, -1, -1, -1)$
4B	0	—
3A	4	$(\omega, \omega, \omega^2, \omega^2)$
6A	0	—
8A	0	—
8B	0	—
7A	3	$(\zeta^{3^j}, \zeta^{3^j \cdot 2}, \zeta^{3^j \cdot 4})$
7B	3	$(\zeta^{3^j-1}, \zeta^{3^j-1 \cdot 2}, \zeta^{3^j-1 \cdot 4})$

Here $\omega = e^{2i\pi/3}$.

Table 5: Decomposition of $H^0(X_1, K_{X_1}^d)$

d	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$a_1(d)$	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	1	0	1
$a_2(d)$	1	0	0	0	0	0	1	1	1	0	1	0	1	1	2	1	1	1	1	1
$a_3(d)$	0	0	1	0	1	0	0	0	1	1	1	1	1	0	1	1	2	1	2	1
$a_4(d)$	0	1	0	1	0	1	0	2	1	2	1	2	1	2	2	3	2	3	2	3
$a_5(d)$	0	0	1	0	1	1	1	1	2	1	2	2	2	2	3	2	3	3	3	3
$a_6(d)$	0	0	1	0	1	1	1	1	2	1	2	2	2	2	3	2	3	3	3	3
$a_7(d)$	0	0	1	0	1	1	1	1	2	1	2	2	2	2	3	2	3	3	3	3
$a_8(d)$	0	0	0	1	1	1	2	1	1	2	2	2	3	3	2	3	3	3	4	4
$a_9(d)$	1	0	1	1	2	1	3	2	3	3	4	3	5	4	5	5	6	5	7	6
$a_{10}(d)$	0	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10
$a_{11}(d)$	0	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10

 Table 6: Decomposition of $H^0(X_2, K_{X_2}^d)$

d	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$a_1(d)$	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	1	0	1
$a_2(d)$	0	0	1	0	1	0	0	0	1	1	1	1	1	0	1	1	2	1	2	1
$a_3(d)$	1	0	0	0	0	0	1	1	1	0	1	0	1	1	2	1	1	1	1	1
$a_4(d)$	0	1	0	1	0	1	0	2	1	2	1	2	1	2	2	3	2	3	2	3
$a_5(d)$	0	0	1	0	1	1	1	1	2	1	2	2	2	2	3	2	3	3	3	3
$a_6(d)$	0	0	1	0	1	1	1	1	2	1	2	2	2	2	3	2	3	3	3	3
$a_7(d)$	0	0	1	0	1	1	1	1	2	1	2	2	2	2	3	2	3	3	3	3
$a_8(d)$	0	0	0	1	1	1	2	1	1	2	2	2	3	3	2	3	3	3	4	4
$a_9(d)$	1	0	1	1	2	1	3	2	3	3	4	3	5	4	5	5	6	5	7	6
$a_{10}(d)$	0	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10
$a_{11}(d)$	0	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10

5 The quadratic equations and the matrix representation of G

From Tables 5 and 6 we have

$$H^0(X_i, K_{X_i}) = W_9 \oplus W_{i+1}, \quad i=1,2, \quad (7)$$

and

$$H^0(X_i, K_{X_i}^{\otimes 2}) = W_4 \oplus W_{10} \oplus W_{11}. \quad (8)$$

Table 7: Decomposition of $S^d(W_9 + W_2)$

d	0	1	2	3	4	5	6	7	8	9	10
$a_1(d)$	1	0	1	0	9	5	85	137	637	1385	4210
$a_2(d)$	0	1	0	4	6	57	145	594	1565	4716	11616
$a_3(d)$	0	0	0	5	6	57	145	594	1562	4716	11619
$a_4(d)$	0	0	3	1	34	71	387	1010	3473	8843	24274
$a_5(d)$	0	0	0	8	20	118	374	1324	3761	10814	27442
$a_6(d)$	0	0	0	8	19	117	367	1317	3743	10796	27391
$a_7(d)$	0	0	0	8	19	117	367	1317	3743	10796	27391
$a_8(d)$	0	0	1	5	32	123	447	1470	4398	12171	31680
$a_9(d)$	0	1	0	13	41	234	725	2646	7486	21564	54810
$a_{10}(d)$	0	0	3	13	80	306	1187	3783	11569	31765	83234
$a_{11}(d)$	0	0	3	13	80	306	1187	3783	11569	31765	83234

Table 8: Decomposition of $S^d(W_9 + W_3)$

d	0	1	2	3	4	5	6	7	8	9	10
$a_1(d)$	1	0	1	0	9	5	85	137	637	1385	4210
$a_2(d)$	0	0	0	5	6	57	145	594	1562	4716	11619
$a_3(d)$	0	1	0	4	6	57	145	594	1565	4716	11616
$a_4(d)$	0	0	3	1	34	71	387	1010	3473	8843	24274
$a_5(d)$	0	0	0	8	20	118	374	1324	3761	10814	27442
$a_6(d)$	0	0	0	8	19	117	367	1317	3743	10796	27391
$a_7(d)$	0	0	0	8	19	117	367	1317	3743	10796	27391
$a_8(d)$	0	0	1	5	32	123	447	1470	4398	12171	31680
$a_9(d)$	0	1	0	13	41	234	725	2646	7486	21564	54810
$a_{10}(d)$	0	0	3	13	80	306	1187	3783	11569	31765	83234
$a_{11}(d)$	0	0	3	13	80	306	1187	3783	11569	31765	83234

Table 9: Decomposition of $S^d W_9$

d	0	1	2	3	4	5	6	7	8	9	10
1	1	0	1	0	4	2	34	40	183	332	922
2	0	0	0	2	3	22	53	184	433	1140	2501
3	0	0	0	2	3	22	53	184	433	1140	2501
4	0	0	2	0	18	28	146	308	980	2120	5284
5	0	0	0	4	10	46	140	408	1047	2612	5922
6	0	0	0	5	9	50	133	417	1030	2643	5878
7	0	0	0	5	9	50	133	417	1030	2643	5878
8	0	0	1	2	17	48	165	454	1230	2928	6863
9	0	1	0	7	20	100	260	843	2060	5267	11772
10	0	0	2	8	39	130	431	1200	3200	7740	17928
11	0	0	2	8	39	130	431	1200	3200	7740	17928

So we assume that $X_i \subset \mathbb{P}^{16} = \mathbb{P}(V_i^*)$, where V_i has the structure of the representation $W_9 \oplus W_{i+1}$. From Tables 7 and 8 we have

$$S^2(V_i) = W_1 \oplus W_4^3 \oplus W_8 \oplus W_{10}^3 \oplus W_{11}^3, \quad i=1,2. \quad (9)$$

Table 10: Decomposition of $S^d W_2$

d	0	1	2	3	4	5	6	7	8	9	10
$a_1(d)$	1	0	0	0	1	0	1	0	1	0	1
$a_2(d)$	0	1	0	0	0	1	0	1	1	2	0
$a_3(d)$	0	0	0	1	0	1	0	1	0	2	1
$a_4(d)$	0	0	1	0	1	0	2	0	3	1	4
$a_5(d)$	0	0	0	1	0	1	1	2	1	3	2
$a_6(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_7(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_8(d)$	0	0	0	0	1	1	1	2	2	2	3
$a_9(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_{10}(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_{11}(d)$	0	0	0	0	0	0	0	0	0	0	0

Table 11: Decomposition of $S^d W_3$

d	0	1	2	3	4	5	6	7	8	9	10
$a_1(d)$	1	0	0	0	1	0	1	0	1	0	1
$a_2(d)$	0	0	0	1	0	1	0	1	0	2	1
$a_3(d)$	0	1	0	0	0	1	0	1	1	2	0
$a_4(d)$	0	0	1	0	1	0	2	0	3	1	4
$a_5(d)$	0	0	0	1	0	1	1	2	1	3	2
$a_6(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_7(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_8(d)$	0	0	0	0	1	1	1	2	2	2	3
$a_9(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_{10}(d)$	0	0	0	0	0	0	0	0	0	0	0
$a_{11}(d)$	0	0	0	0	0	0	0	0	0	0	0

It follows that

$$H^0(\mathbb{P}^{16}, \mathcal{I}_{X_i}(2)) = W_1 \oplus W_4^2 \oplus W_8 \oplus W_{10}^2 \oplus W_{11}^2, \quad i=1,2. \quad (10)$$

Where $\mathcal{I}_{X_i}$ denotes the ideal sheaf of X_i in $\mathbb{P}^{16}$. From now on we will use the notation

$$I_{X_i}(2) = H^0(\mathbb{P}^{16}, \mathcal{I}_{X_i}(2)).$$

We have

$$\pi_{W_1}(I_{X_i}(2)) = \pi_{W_1}(S^2 V_i), \quad (11)$$

and

$$\pi_{W_8}(I_{X_i}(2)) = \pi_{W_8}(S^2 V_i). \quad (12)$$

The remaining spaces $\pi_{W_4}(I_{X_i}(2))$, $\pi_{W_{10}}(I_{X_i}(2))$ and $\pi_{W_{11}}(I_{X_i}(2))$ will be found in Sections 6,7 and 8 below.

For calculations we fix an action of G on $\mathbb{P}^{16}$. In V_i (respectively V_i^*) we consider a basis $\{y_1, \dots, y_{17}\}$ (respectively the dual basis $\{e_1, \dots, e_{17}\}$). Sometimes we use the following subspaces of V_i

$$V_{i,9} = \langle y_1, \dots, y_{14} \rangle, \quad (13)$$

$$V_{i,i+1} = \langle y_{15}, y_{16}, y_{17} \rangle. \quad (14)$$

With respect to these bases, the action of the generators P and Q on V_i (respectively V_i^*) is given by the matrices $\text{diag}(P_{W_9}, P_{W_{i+1}})$ and $\text{diag}(Q_{W_9}, Q_{W_{i+1}})$ (respectively the inverse transpose of these matrices), where the matrices $P_{W_j}, Q_{W_j}, j = 9, 2$ are defined in formulae (15), (16), (17) and (18) (recall that $\omega = e^{2i\pi/3}$ and $\zeta = e^{2i\pi/7}$). The matrices P_{W_3}, Q_{W_3} could be taken as the respective inverse transpose of P_{W_2}, Q_{W_2} however we will focus only on the curve X_1 .

Remark 5.1. One can identify G with a subgroup of $\text{Aut}(G)$. The action of G extends to an action of $\text{Aut}(G)$ on V_1 . So the exterior automorphisms move $X_1 \subset \mathbb{P}(V_1^*)$ giving rise to another copy $X'_1 \subset \mathbb{P}(V_1^*)$ of X_1 . For instance, we can assume $\text{Aut}(G) = \langle P, Q, E \rangle$, where E is such that the action on V_1 is given by the matrix $\text{diag}(E_{W_9}, E_{W_2})$, where E_{W_9} and E_{W_2} are defined in formulae (19) and (21) respectively.

Remark 5.2. Let Y denote the Klein quartic. We have that Y is isomorphic to the quotient X_i/N for $i = 1, 2$, where N is the normal subgroup of G generated by the involutions in the class $2A$. Let $f_i : X_i \mapsto Y$ be the quotient maps. Then we have that the action of G induced on Y from X_1 or X_2 is such that $d_Y h_{7B} = (\zeta^1, \zeta^2, \zeta^4)$ or $d_Y h_{7B} = (\zeta^3, \zeta^5, \zeta^6)$ respectively. The action induced by f_1 or f_2 gives $H^0(Y, K_Y)$ the structure of the representation W_2 or W_3 respectively. Since $f_i^* K_Y \cong K_{X_i}$, we see that f_i can be recovered by composing the canonical embedding $X_i \subset \mathbb{P}(V_i^*)$ with the projection $\pi_i : \mathbb{P}(V_i^*) \dashrightarrow \mathbb{P}(V_{i,i+1}^*)$.

$$P_{W_2} = \begin{pmatrix} \zeta^3 & 0 & 0 \\ 0 & \zeta^5 & 0 \\ 0 & 0 & \zeta^6 \end{pmatrix} \quad (15)$$

$$Q_{W_2} = \frac{1}{\sqrt{-7}} \begin{pmatrix} \zeta^4 - \zeta^6 & -1 + \zeta^3 & \zeta - \zeta^2 \\ -\zeta + \zeta^4 & \zeta^2 - \zeta^3 & -1 + \zeta^5 \\ -1 + \zeta^6 & \zeta^2 - \zeta^4 & \zeta - \zeta^5 \end{pmatrix} \quad (16)$$

(17)

(18)

(19)

(21)

6 The generators of $\pi_{\mathbf{W}_4}(I_{X_1}(2))$

Let $h_{7B} = P^2QP^3Q = (1, 8, 5, 4, 11, 2, 13)(3, 12, 9, 7, 14, 6, 10)$ be the representative of the conjugacy class $7B$ of G . From Table 3 we see that $\rho(h_{7B})_{\zeta^2}(\mathbf{W}_4)$ has dimension 1. So, since $\mathbf{W}_4$ is an irreducible $\mathbb{C}G$ -module, the subspace $\rho(h_{7B})_{\zeta^2}(\pi_{\mathbf{W}_4}(I_{X_1}(2)))$ generates $\pi_{\mathbf{W}_4}(I_{X_1}(2))$ as a $\mathbb{C}G$ -module. We shall find a point $p \in X_1$ such that not all ζ^2 -eigenquadrics of h_{7B} in $\pi_{\mathbf{W}_4}(S^2\mathbf{V}_1)$ vanish on p . Then the subspace $\rho(h_{7B})_{\zeta^2}(\pi_{\mathbf{W}_4}(I_{X_1}(2)))$ is the set of all the ζ^2 -eigenquadrics of h_{7B} in $\pi_{\mathbf{W}_4}(S^2\mathbf{V}_1)$ that vanish on p .

If p is a fixed point of h_{7B} in X_1 , then it lives in a subspace $\mathbb{P}(\rho(h_{7B})_{\zeta^i}(\mathbf{V}_1^*)) \subset \mathbb{P}(\mathbf{V}_1^*)$ for some i . It can be checked that $i = 6, 5$ or 3 if $dh_{7B}p = \zeta^1, \zeta^2$ or ζ^4 respectively.

Now we will look for the fixed point p of h_{7B} in X_1 lying on $\mathbb{P}(\rho(h_{7B})_{\zeta^6}(\mathbf{V}_1^*))$. Note that $\rho(h_{7B})_{\zeta^6}(\mathbf{V}_{1,9}^*)$ has dimension 2 (see Table 3) and that $f_1(p)$ (see Remark 5.2) lies on the subspace $\rho(h_{7B})_{\zeta^6}(\mathbf{V}_{1,2}^*)$ which is one dimensional. Therefore we can assume that

$$p = \alpha \cdot v_1 + \beta \cdot v_2 + v_3, \quad (22)$$

where $\alpha, \beta \in \mathbb{C}$ and $\{v_1, v_2\}$ and $\{v_3\}$ are bases for $\rho(h_{7B})_{\zeta^6}(\mathbf{V}_{1,9}^*)$ and $\rho(h_{7B})_{\zeta^6}(\mathbf{V}_{1,2}^*)$ respectively.

To find α and β we will evaluate on the point p the ζ^2 -eigenquadric q_{ζ^2} of h_{7B} in $\pi_{\mathbf{W}_8}(I_{X_1}(2))$; namely

$$q_{\zeta^2} = \rho(h_{7B})_{\zeta^2}(\pi_{\mathbf{W}_8}(y_1^2)). \quad (23)$$

We take

$$\begin{aligned} v_i &= \rho(h_{7B})_{\zeta^6}(e_i), \quad i = 1, 2 \\ &\text{and} \\ v_3 &= \rho(h_{7B})_{\zeta^6}(e_{17}). \end{aligned} \quad (24)$$

We see that

$$q_{\zeta^2}(v_i) \neq 0, \quad i = 1, 2. \quad (25)$$

We assume that $\beta = 1$ since we always can move the fixed point p by a G -equivariant change of coordinates $t_\lambda : e_i \rightarrow \lambda e_i, i = 1, \dots, 14$ and $e_i \rightarrow e_i, i = 15, 16, 17$. So we have

$$\alpha^2 + 1/7(-6\xi - 4\xi^2 + 4\xi^4 - 3\xi^5 - 2\xi^8 + 3\xi^{10} + 6\xi^{11} - \xi^{13} + 2\xi^{16} + \xi^{17} - 2\xi^{19} + 2\xi^{20}) = 0, \quad (26)$$

where $\xi = e^{2i\pi/21}$.

Notice that the corresponding points to each root of (26) live in projectively equivalent curves in $\mathbb{P}(V_1^*)$. One is obtained from the other by applying the exterior automorphism in Remark 5.1 followed by the equivariant translation t_λ with suitable λ .

Now, a basis for the ζ^2 -eigenspace of $h_{7B}|_{\pi_{W_4}(S^2V_1)}$ is given by $\{b_1, b_2, b_3\}$, where

$$\begin{aligned} b_1 &= \rho(h_{7B})_{\zeta^2}(\pi_{W_4}(y_1^2)), \\ b_2 &= \rho(h_{7B})_{\zeta^2}(\pi_{W_4}(y_1y_2)), \\ b_3 &= y_{17}^2. \end{aligned} \tag{27}$$

Evaluating on p one proves that $b_i(p) \neq 0$, $i = 1, 2, 3$. Thus the quadrics

$$\vartheta_1 = b_1(p)b_2 - b_2(p)b_1 \text{ and } \vartheta_2 = b_1(p)b_3 - b_3(p)b_1 \tag{28}$$

form a basis for the ζ^2 -eigenspace of $h_{7B}|_{\pi_{W_4}(I_{X_1}(2))}$.

7 The generators of $\pi_{W_{10}}(I_{X_1}(2))$

We shall find linearly independent elements $\theta_1, \theta_2, \theta_3 \in \pi_{W_{10}}(S^2V_1)$ such that the G -module generated by each one of them is isomorphic to W_{10} and such that $\theta_1, \theta_2, \theta_3$ are G -equivalent, meaning that if ϕ_{ij} is an isomorphism of G -modules from the G -module generated by θ_i to that generated by θ_j then $\phi_{ij}(\theta_i) = \lambda_{ij}\theta_j$ for some $\lambda_{ij} \in \mathbb{C} \setminus \{0\}$. We take θ_i to be a ζ^2 -eigenquadric of $h_{7B}|_{\pi_{W_{10}}(S^2V_1)}$. Let p be the fixed point of h_{7B} we found in Section 6. If $\theta_i(p) \neq 0$, $i = 1, 2, 3$ then

$$\begin{aligned} \wp_1 &= \theta_1(p)\theta_2 - \theta_2(p)\theta_1, \\ \wp_2 &= \theta_1(p)\theta_3 - \theta_3(p)\theta_1 \end{aligned} \tag{29}$$

generate $\pi_{W_{10}}(I_{X_1}(2))$.

We first notice that

$$S^2V_1 = S^2V_{1,9} \oplus S^2V_{1,2} \oplus V_{1,9} \otimes V_{1,2}. \tag{30}$$

Using that (see Tables 9,10)

$$\begin{aligned} S^2V_{1,9} &= W_1 \oplus W_4^2 \oplus W_8 \oplus W_{10}^2 \oplus W_{11}^2, \\ S^2V_{1,2} &= W_4, \end{aligned} \tag{31}$$

one concludes that

$$V_{1,9} \otimes V_{1,2} = W_{10} \oplus W_{11}. \quad (32)$$

Then we define an isomorphism of G -modules

$$\varrho_{W_{10}} : \pi_{W_{10}}(V_{1,9} \otimes V_{1,2}) \oplus \pi_{W_{10}}(V_{1,9} \otimes V_{1,2}) \rightarrow \pi_{W_{10}}(S^2 V_{1,9}).$$

So given

$$\theta_1 \in \pi_{W_{10}}(V_{1,9} \otimes V_{1,2})$$

one can take

$$\begin{aligned} \theta_2 &= \varrho_{W_{10}}((\theta_1, 0)) \\ \theta_3 &= \varrho_{W_{10}}((0, \theta_1)). \end{aligned} \quad (33)$$

To construct $\varrho_{W_{10}}$ we consider the bases B_1 of $\pi_{W_{10}}(V_{1,9} \otimes V_{1,2})$ and B_2 of $\pi_{W_{10}}(S^2 V_{1,9})$ given in formulae (35) and (37). Let A, C be the matrix representations of G corresponding to the action on $\pi_{W_{10}}(V_{1,9} \otimes V_{1,2})$ and on $\pi_{W_{10}}(S^2 V_{1,9})$ with respect to these bases. If M_0 is a 42×42 complex matrix such that $\det S \neq 0$, where S given by equation (34), then one can take $\varrho_{W_{10}}$ to be the linear map induced by S with respect to the bases $B_1 \cup B_2$ and B_2 .

$$S = \sum_{h \in G} C_h \cdot M_0 \cdot \text{diag}(A_h, A_h)^{-1}. \quad (34)$$

For our calculation we use $M_0 = I_{42}$ the identity matrix.

$$B_1 = \{\mathbb{k}_{1,j}, \mathbb{k}_{2,j}, \mathbb{k}_{3,j} \mid j = 1 \dots 7\}, \quad (35)$$

where

$$\begin{aligned} \mathbb{k}_{i,j} &= \rho(h_{7B})_{\zeta^j}(Q \cdot \mathbb{k}_{i,2}), \text{ for } j \neq 2, \\ \mathbb{k}_{1,2} &= \rho(h_{7B})_{\zeta^2}(\pi_{W_{10}}(y_1 y_{15})), \\ \mathbb{k}_{i+1,2} &= \rho(h_{7B})_{\zeta^2}(Q \cdot \mathbb{k}_{i,2}). \end{aligned} \quad (36)$$

$$B_2 = \{\kappa_{1,j}, \kappa_{2,j}, \kappa_{3,j}, \kappa_{4,j}, \kappa_{5,j}, \kappa_{6,j} \mid j = 1 \dots 7\}, \quad (37)$$

where

$$\begin{aligned} \kappa_{i,j} &= \rho(h_{7B})_{\zeta^j}(Q \cdot \kappa_{i,2}), \text{ for } j \neq 2, \\ \kappa_{1,2} &= \rho(h_{7B})_{\zeta^2}(\pi_{W_{10}}(y_1 y_3)), \\ \kappa_{4,2} &= \rho(h_{7B})_{\zeta^2}(Q \cdot \rho(h_{7B})_{\zeta^3}(Q \cdot \kappa_{1,2})), \\ \kappa_{i+1,2} &= \rho(h_{7B})_{\zeta^2}(Q \cdot \kappa_{i,2}), \text{ for } i = 1, 2, 4, 5. \end{aligned} \quad (38)$$

We take $\theta_1 = \mathbb{k}_{1,2}$. Then

$$\begin{aligned} \theta_2 &= \sum_{i=1}^6 s_{i+6,4} \kappa_{i,2}, \\ \theta_3 &= \sum_{i=1}^6 s_{i+6,25} \kappa_{i,2}. \end{aligned} \quad (39)$$

8 The generators of $\pi_{\mathbf{w}_{11}}(I_{X_1}(2))$

The situation is similar to that of $\pi_{\mathbf{w}_{10}}(I_{X_1}(2))$. We define an isomorphism of G -modules

$$\varrho_{\mathbf{w}_{11}} : \pi_{\mathbf{w}_{11}}(\mathbf{V}_{1,9} \otimes \mathbf{V}_{1,2}) \oplus \pi_{\mathbf{w}_{11}}(\mathbf{V}_{1,9} \otimes \mathbf{V}_{1,2}) \rightarrow \pi_{\mathbf{w}_{11}}(S^2 \mathbf{V}_{1,9}).$$

which corresponds to a matrix

$$S' = \sum_{h \in G} C'_h \cdot M'_0 \cdot \text{diag}(A'_h, A'_h)^{-1}. \quad (40)$$

where A', C' are the matrix representations of G corresponding to the action on $\pi_{\mathbf{w}_{11}}(\mathbf{V}_{1,9} \otimes \mathbf{V}_{1,2})$ and on $\pi_{\mathbf{w}_{11}}(S^2 \mathbf{V}_{1,9})$ with respect to fixed bases B'_1 and B'_2 respectively (see formulae (41), (43)). We use $M'_0 = I_{42}$ the identity matrix. Then $\wp'_1$ and $\wp'_2$ in formula (46) generate $\pi_{\mathbf{w}_{11}}(I_{X_1}(2))$.

$$B'_1 = \{\mathbb{k}'_{1,j}, \mathbb{k}'_{2,j}, \mathbb{k}'_{3,j} \mid j = 1 \dots 7\}, \quad (41)$$

where

$$\begin{aligned} \mathbb{k}'_{i,j} &= \rho(h_{7B})_{\zeta^j}(Q \cdot \mathbb{k}'_{i,2}), \text{ for } j \neq 2, \\ \mathbb{k}'_{1,2} &= \rho(h_{7B})_{\zeta^2}(\pi_{\mathbf{w}_{11}}(y_1 y_{15})), \\ \mathbb{k}'_{i+1,2} &= \rho(h_{7B})_{\zeta^2}(Q \cdot \mathbb{k}'_{i,2}). \end{aligned} \quad (42)$$

$$B'_2 = \{\kappa'_{1,j}, \kappa'_{2,j}, \kappa'_{3,j}, \kappa'_{4,j}, \kappa'_{5,j}, \kappa'_{6,j} \mid j = 1 \dots 7\}, \quad (43)$$

where

$$\begin{aligned} \kappa'_{i,j} &= \rho(h_{7B})_{\zeta^j}(Q \cdot \kappa'_{i,2}), \text{ for } j \neq 2, \\ \kappa'_{1,2} &= \rho(h_{7B})_{\zeta^2}(\pi_{\mathbf{w}_{11}}(y_1 y_3)), \\ \kappa'_{4,2} &= \rho(h_{7B})_{\zeta^2}(Q \cdot \rho(h_{7B})_{\zeta^3}(Q \cdot \kappa'_{1,2})), \\ \kappa'_{i+1,2} &= \rho(h_{7B})_{\zeta^2}(Q \cdot \kappa'_{i,2}) \text{ for } i = 1, 2, 4, 5. \end{aligned} \quad (44)$$

Taking $\theta'_1 = \mathbb{k}'_{1,2}$ we take

$$\begin{aligned} \theta'_2 &= \sum_{i=1}^6 s'_{i+6,4} \kappa'_{i,2}, \\ \theta'_3 &= \sum_{i=1}^6 s'_{i+6,25} \kappa'_{i,2}. \end{aligned} \quad (45)$$

One checks that $\theta'_1(p)\theta'_2(p)\theta'_3(p) \neq 0$. So we define

$$\begin{aligned} \wp'_1 &= \theta'_1(p)\theta'_2 - \theta'_2(p)\theta'_1, \\ \wp'_2 &= \theta'_1(p)\theta'_3 - \theta'_3(p)\theta'_1. \end{aligned} \quad (46)$$

Although one can speed up the calculation of (34) and (40) by using representatives of cosets of a subgroup of G , the computation was still very slow.

9 A basis for $I_{X_1}(2)$

Here we define a basis for $I_{X_1}(2)$ by giving bases D_i , $i = 1, 4, 8, 10, 11$, for the subspaces $\pi_{w_i}(I_{X_1}(2))$. Let α be a solution of (26) and p be the corresponding point. Let q_{ζ^2} be as defined in (23). Let ϑ_1, ϑ_2 be as defined in (28). Let $\wp_1, \wp_2, \wp'_1$ and $\wp'_2$ be as defined in (29) and (46) respectively. Then we have

$$\begin{aligned} D_1 &= \{d_{1,7,1}\}, \\ \text{where} \quad d_{1,7,1} &= \pi_{w_1}(y_1 y_2). \end{aligned} \tag{47}$$

$$\begin{aligned} D_4 &= D_{4,1} \cup D_{4,2}, \\ \text{where} \quad D_{4,i} &= \{d_{4,j,i} \mid j=1, \dots, 6.\}, \quad i=1,2, \\ \text{and} \quad d_{4,j,i} &= \rho(h_{7B})_{\zeta^j}(Q \cdot \vartheta_i). \end{aligned} \tag{48}$$

$$\begin{aligned} D_8 &= \{d_{8,j,1} \mid j=1, \dots, 7.\} \cup \{d_{8,7,2}\}, \\ \text{where} \end{aligned} \tag{49}$$

$$\begin{aligned} d_{8,j,1} &= \rho(h_{7B})_{\zeta^j}(Q \cdot q_{\zeta^2}), \\ d_{8,7,2} &= \rho(h_{7B})_1(Q^2 \cdot q_{\zeta^2}). \end{aligned}$$

$$\begin{aligned} D_{10} &= D_{10,1} \cup D_{10,2}, \\ \text{where} \quad D_{10,i} &= \phi_{\theta_1, \wp_i}(B_1), \\ B_1 &\text{ is defined in formula (35) and} \\ \phi_{\theta_1, \wp_i} &: \mathbb{C}G_{\langle \theta_1 \rangle} \rightarrow \mathbb{C}G_{\langle \wp_i \rangle} \end{aligned} \tag{50}$$

is the isomorphism of G -modules that maps θ_1 to $\wp_i$. We write

$$\begin{aligned} d_{10,j,k} &= \phi_{\theta_1, \wp_1}(\mathbb{k}_{k,j}), \\ d_{10,j,k+3} &= \phi_{\theta_1, \wp_2}(\mathbb{k}_{k,j}), \quad k = 1, 2, 3. \end{aligned}$$

One has for instance

$$\begin{aligned} d_{10,2,1} &= \wp_1, \\ d_{10,2,k+1} &= \rho(h_{7B})_{\zeta^2}(Q \cdot d_{10,2,k}), \quad k=1,2. \\ d_{10,j,k} &= \rho(h_{7B})_{\zeta^j}(Q \cdot d_{10,2,k}), \quad j=1,3,4,5,6,7, \quad k=1,2,3. \end{aligned}$$

Similarly we define

$$D_{11} = D_{11,1} \cup D_{11,2}, \quad (51)$$

where

$$D_{11,i} = \{d_{11,j,3(i-1)+k} \mid j=1, \dots, 7, k=1,2,3\}, i=1,2.$$

and

$$d_{11,j,3(i-1)+k} = \phi_{\theta'_1, \phi'_i}(\mathbb{k}'_{k,j}).$$

10 The space of cubic forms

Our main conclusion is the following theorem.

Theorem 10.1. *The homogeneous ideal of the canonical embedding of X_1 in $\mathbb{P}^{16}$ is generated by the quadratic forms in it.*

For the proof we used the basis defined in Section 9 to show that the space of cubic forms generated by $I_{X_1}(2)$ is $I_{X_1}(3)$, namely $V_1 \cdot I_{X_1}(2) = I_{X_1}(3)$. To do that it is enough to show that $\pi_{W_i}(V_1 \cdot I_{X_1}(2)) = \pi_{W_i}(I_{X_1}(3))$, $i=1, \dots, 11$.

So we computed the dimensions for the eigenspaces $\rho(h_{7B})_{\zeta^j} \left(\pi_{W_i}(V_1 \cdot I_{X_1}(2)) \right)$, for some j . Thus the multiplicity of the representation W_i in $\pi_{W_i}(V_1 \cdot I_{X_1}(2))$ can be deduced from the information in Table 3. In Table 12 we list the dimensions of the spaces $\rho(h_{7B})_{\zeta^j} \left(\pi_{W_i}(V_1 \cdot I_{X_1}(2)) \right)$ and we also list bases for these spaces. On the other hand, from Tables 5 and 7 we have

$$H^0(X_1, K_{X_1}^{\otimes 3}) = W_3 \oplus W_5 \oplus W_6 \oplus W_7 \oplus W_9 \oplus W_{10} \oplus W_{11},$$

and

$$S^3(V_1) = W_2^4 \oplus W_3^5 \oplus W_4 \oplus W_5^8 \oplus W_6^8 \oplus W_7^8 \oplus W_8^5 \oplus W_9^{13} \oplus W_{10}^{13} \oplus W_{11}^{13},$$

then

$$H^0(\mathbb{P}^{16}, \mathcal{I}_{X_1}(3)) = W_2^4 \oplus W_3^4 \oplus W_4 \oplus W_5^7 \oplus W_6^7 \oplus W_7^7 \oplus W_8^5 \oplus W_9^{12} \oplus W_{10}^{12} \oplus W_{11}^{12}. \quad (52)$$

Table 12: Generators for $I_{X_1}(3)$

$W_{i,j}$	dimension of $\rho(h_{7B})_{\zeta^j}(\pi_{\mathbf{w}_i}(\mathbf{V}_1 \cdot I_{X_1}(2)))$	basis for $\rho(h_{7B})_{\zeta^j}(\pi_{\mathbf{w}_i}(\mathbf{V}_1 \cdot I_{X_1}(2)))$
$W_1, -$	0	—
$W_2, 2$	4	$\pi_{\mathbf{w}_2} \left(\left\{ \begin{array}{l} [y_{16}]_{\zeta^2} d_{1,7,1}, [y_{17}]_{\zeta^1} d_{8,1,1}, \\ [y_1]_{\zeta^0} d_{10,2,1}, [y_1]_{\zeta^0} d_{10,2,4} \end{array} \right\} \right)$
$W_3, 3$	4	$\pi_{\mathbf{w}_3} \left(\left\{ \begin{array}{l} [y_{17}]_{\zeta^1} d_{4,2,1}, [y_{17}]_{\zeta^1} d_{4,2,2}, \\ [y_1]_{\zeta^1} d_{10,2,1}, [y_1]_{\zeta^1} d_{10,2,4} \end{array} \right\} \right)$
$W_4, 2$	1	$\pi_{\mathbf{w}_4} \left(\left\{ [y_{17}]_{\zeta^1} d_{8,1,1} \right\} \right)$
$W_5, 2$	7	$\pi_{\mathbf{w}_5} \left(\left\{ \begin{array}{l} [y_{17}]_{\zeta^1} d_{4,1,1}, [y_{17}]_{\zeta^1} d_{4,1,2}, [y_{17}]_{\zeta^1} d_{8,1,1}, \\ [y_1]_{\zeta^1} d_{10,1,1}, [y_1]_{\zeta^1} d_{10,1,2}, [y_1]_{\zeta^1} d_{10,1,4}, \\ [y_1]_{\zeta^1} d_{11,1,1} \end{array} \right\} \right)$
$W_6, 2$	7	$\pi_{\mathbf{w}_6} \left(\left\{ \begin{array}{l} [y_1]_{\zeta^1} d_{8,1,1}, [y_1]_{\zeta^1} d_{4,1,1}, [y_1]_{\zeta^1} d_{4,1,2}, \\ [y_1]_{\zeta^1} d_{10,1,1}, [y_1]_{\zeta^1} d_{10,1,2}, [y_1]_{\zeta^1} d_{10,1,4}, \\ [y_1]_{\zeta^1} d_{10,1,5} \end{array} \right\} \right)$
$W_7, 2$	7	$\pi_{\mathbf{w}_7} \left(\left\{ \begin{array}{l} [y_1]_{\zeta^1} d_{8,1,1}, [y_1]_{\zeta^1} d_{4,1,1}, [y_1]_{\zeta^1} d_{4,1,2}, \\ [y_1]_{\zeta^1} d_{10,1,1}, [y_1]_{\zeta^1} d_{10,1,4}, [y_{17}]_{\zeta^1} d_{10,1,1}, \\ [y_{17}]_{\zeta^1} d_{10,1,4} \end{array} \right\} \right)$
$W_8, 2$	5	$\pi_{\mathbf{w}_8} \left(\left\{ \begin{array}{l} [y_1]_{\zeta^1} d_{10,1,1}, [y_1]_{\zeta^1} d_{10,1,2}, [y_1]_{\zeta^1} d_{10,1,4}, \\ [y_1]_{\zeta^1} d_{11,1,1}, [y_1]_{\zeta^1} d_{11,1,2} \end{array} \right\} \right)$
$W_9, 2$	24	$\pi_{\mathbf{w}_9} \left(\left\{ \begin{array}{l} [y_1]_{\zeta^1} [D_4]_{\zeta^1} \cup [y_2]_{\zeta^1} [D_4]_{\zeta^1} \cup [y_1]_{\zeta^3} [D_4]_{\zeta^6} \cup \\ [y_2]_{\zeta^3} [D_4]_{\zeta^6} \cup [y_1]_{\zeta^1} [D_{10}]_{\zeta^1} \cup \\ [y_1]_{\zeta^1} d_{8,1,1}, [y_2]_{\zeta^1} d_{8,1,1}, [y_2]_{\zeta^1} d_{10,1,1}, \\ [y_2]_{\zeta^1} d_{10,1,2}, [y_{17}]_{\zeta^1} d_{10,1,1}, [y_{17}]_{\zeta^1} d_{10,1,2}, \\ [y_1]_{\zeta^2} d_{10,7,1}, [y_2]_{\zeta^1} d_{10,1,4}, [y_2]_{\zeta^1} d_{10,1,5} \\ [y_{17}]_{\zeta^1} d_{10,1,4} \end{array} \right\} \right)$
$W_{10}, 2$	36	$\pi_{\mathbf{w}_{10}} \left(\left\{ \begin{array}{l} [y_1]_{\zeta^1} [D_4]_{\zeta^1} \cup [y_2]_{\zeta^1} [D_4]_{\zeta^1} \cup [y_1]_{\zeta^3} [D_4]_{\zeta^6} \cup \\ [y_1]_{\zeta^2} [D_8]_{\zeta^7} \cup [y_2]_{\zeta^2} [D_8]_{\zeta^7} \cup [y_{17}]_{\zeta^1} [D_{11}]_{\zeta^1} \cup \\ [y_1]_{\zeta^1} [D_{10,1}]_{\zeta^1} \cup [y_2]_{\zeta^1} [D_{10,1}]_{\zeta^1} \cup \\ [y_1]_{\zeta^2} [D_{10,1}]_{\zeta^7} \cup [y_2]_{\zeta^2} [D_{10,1}]_{\zeta^7} \cup \\ [y_1]_{\zeta^1} d_{8,1,1}, [y_2]_{\zeta^1} d_{8,1,1}, [y_{17}]_{\zeta^1} d_{10,1,1}, \\ [y_{17}]_{\zeta^1} d_{10,1,2}, [y_{16}]_{\zeta^2} d_{10,7,1}, [y_{17}]_{\zeta^1} d_{10,1,4}, \\ [y_{17}]_{\zeta^1} d_{10,1,5}, [y_{16}]_{\zeta^2} d_{10,7,4} \end{array} \right\} \right)$
$W_{11}, 2$	36	$\pi_{\mathbf{w}_{11}} \left(\left\{ \begin{array}{l} [y_1]_{\zeta^1} [D_4]_{\zeta^1} \cup [y_2]_{\zeta^1} [D_4]_{\zeta^1} \cup [y_1]_{\zeta^3} [D_4]_{\zeta^6} \cup \\ [y_1]_{\zeta^2} [D_8]_{\zeta^7} \cup [y_2]_{\zeta^2} [D_8]_{\zeta^7} \cup [y_{17}]_{\zeta^1} [D_{10}]_{\zeta^1} \cup \\ [y_1]_{\zeta^1} [D_{10,1}]_{\zeta^1} \cup [y_2]_{\zeta^1} [D_{10,1}]_{\zeta^1} \cup \\ [y_1]_{\zeta^2} [D_{10,1}]_{\zeta^7} \cup [y_2]_{\zeta^2} [D_{10,1}]_{\zeta^7} \cup \\ [y_1]_{\zeta^1} d_{8,1,1}, [y_2]_{\zeta^1} d_{8,1,1}, [y_{17}]_{\zeta^1} d_{11,1,1}, \\ [y_{17}]_{\zeta^1} d_{11,1,2}, [y_{16}]_{\zeta^2} d_{11,7,2}, [y_{17}]_{\zeta^1} d_{11,1,4}, \\ [y_{17}]_{\zeta^1} d_{11,1,5}, [y_{16}]_{\zeta^2} d_{11,7,4} \end{array} \right\} \right)$

Here we write $[y_r]_{\zeta^j} = \rho(h_{7B})_{\zeta^j}(y_r)$ and $[D*]_{\zeta^v} = \{d_{i,j,k} \in D* \mid j \equiv v \pmod{7}\}$.

11 The genus 14 case

Let X_1 be the Hurwitz curve of genus 14 considered in [7]. Using the quadrics in the Theorem 2.5 of [7] one also proves the following.

Theorem 11.1. *The homogeneous ideal of the canonical embedding of X_1 in $\mathbb{P}^{13}$ is generated quadrics.*

We proceed exactly as in the proof of Theorem 10.1. In this case we used the Table 13 below, the Table 3 in [7] and the decomposition of $H^0(\mathbb{P}^{13}, \mathcal{I}_{X_1}(3))$ into a sum of irreducible representations of $PSL(2, \mathbb{F}_{13})$. That is (use Tables 5 and 8 in [7]),

$$H^0(\mathbb{P}^{13}, \mathcal{I}_{X_1}(3)) = W_2^5 \oplus W_3^5 \oplus W_4^5 \oplus W_5^6 \oplus W_6^5 \oplus W_7^5 \oplus W_8^4 \oplus W_9^8. \quad (53)$$

When computing the quadrics in Theorem 2.5 of [7] one should have into account two misprints: the first in the definition of ϵ_i (before formula (9)) where ϵ_2 should be equal to $\rho(h_{7A})_{\zeta^1}(e_3)$ and the second in the definition of q (also before formula (9)) where $q = \rho(h_{7A})_{\zeta^5}(\pi_{W_6}(y_1^2))$ should be $q = \rho(h_{7A})_{\zeta^5}(\pi_{W_6}(y_2^2))$.

Table 13: Generators for $I_{X_1}(3)$ in the genus 14 case.

$W_{i,j}$	dimension of $\rho(h_{13A})_{\nu^j}(\pi_{\mathbf{w}_i}(\mathbf{V}_1 \cdot I_{X_1}(2)))$	basis for $\rho(h_{13A})_{\nu^j}(\pi_{\mathbf{w}_i}(\mathbf{V}_1 \cdot I_{X_1}(2)))$
$W_1, -$	0	—
$W_2, 2$	5	$\pi_{\mathbf{w}_2} \left(\left\{ \begin{array}{l} [y_1]_{\nu^3} d_{5,12,1}, [y_1]_{\nu^1} d_{6,1,1}, \\ [y_1]_{\nu^1} d_{7,1,1}, [y_1]_{\nu^1} d_{8,1,1}, [y_1]_{\nu^1} d_{8,1,3} \end{array} \right\} \right)$
$W_3, 1$	5	$\pi_{\mathbf{w}_3} \left(\left\{ \begin{array}{l} [y_1]_{\nu^2} d_{5,12,1}, [y_1]_{\nu^2} d_{6,12,1}, \\ [y_1]_{\nu^1} d_{7,13,1}, [y_1]_{\nu^1} d_{8,13,1}, [y_1]_{\nu^1} d_{8,13,2} \end{array} \right\} \right)$
$W_4, 1$	5	$\pi_{\mathbf{w}_4} \left(\left\{ \begin{array}{l} [y_1]_{\nu^2} d_{5,12,1}, [y_1]_{\nu^3} d_{5,11,1}, \\ [y_1]_{\nu^2} d_{6,12,1}, [y_1]_{\nu^3} d_{6,11,1}, [y_1]_{\nu^1} d_{7,13,1} \end{array} \right\} \right)$
$W_5, 1$	6	$\pi_{\mathbf{w}_5} \left(\left\{ \begin{array}{l} [y_1]_{\nu^3} d_{5,11,1}, [y_1]_{\nu^4} d_{5,10,1}, [y_1]_{\nu^2} d_{6,12,1}, \\ [y_1]_{\nu^3} d_{6,11,1}, [y_1]_{\nu^1} d_{7,13,1}, [y_1]_{\nu^2} d_{7,12,1} \end{array} \right\} \right)$
$W_6, 1$	5	$\pi_{\mathbf{w}_6} \left(\left\{ \begin{array}{l} [y_1]_{\nu^2} d_{5,12,1}, [y_1]_{\nu^3} d_{5,11,1}, [y_1]_{\nu^3} d_{6,11,1}, \\ [y_1]_{\nu^4} d_{6,10,1}, [y_1]_{\nu^1} d_{7,13,1} \end{array} \right\} \right)$
$W_7, 1$	5	$\pi_{\mathbf{w}_7} \left(\left\{ \begin{array}{l} [y_1]_{\nu^2} d_{5,12,1}, [y_1]_{\nu^3} d_{5,11,1}, [y_1]_{\nu^2} d_{6,12,1}, \\ [y_1]_{\nu^3} d_{6,11,1}, [y_1]_{\nu^1} d_{7,13,1} \end{array} \right\} \right)$
$W_8, 1$	4	$\pi_{\mathbf{w}_8} \left(\left\{ \begin{array}{l} [y_1]_{\nu^2} d_{5,12,1}, [y_1]_{\nu^3} d_{5,11,1}, [y_1]_{\nu^2} d_{6,12,1}, \\ [y_1]_{\nu^3} d_{6,11,1} \end{array} \right\} \right)$
$W_9, 1$	8	$\pi_{\mathbf{w}_9} \left(\left\{ \begin{array}{l} [y_1]_{\nu^1} d_{8,13,1}, [y_1]_{\nu^1} d_{8,13,2}, [y_1]_{\nu^2} d_{8,12,1}, \\ [y_1]_{\nu^3} d_{7,11,1}, [y_1]_{\nu^2} d_{5,12,1}, [y_1]_{\nu^3} d_{5,11,1}, \\ [y_1]_{\nu^2} d_{6,12,1}, [y_1]_{\nu^3} d_{6,11,1} \end{array} \right\} \right)$

Using the notation of [7] we write $\mathbf{V}_1 = \langle y_1, \dots, y_{14} \rangle$,
 $[y_r]_{\nu^j} = \rho(h_{13A})_{\nu^j}(y_r)$, $\nu = e^{2i\pi/13}$,
 $d_{i,j,1} = \rho(h_{13A})_{\nu^j}(\pi_{\mathbf{w}_5}(y_2^2))$ for $i = 5, 6$ and $j = 1 \dots 12$,
 $d_{7,j,1} = \rho(h_{13A})_{\nu^j}(h_{7A} \cdot q_0)$ for $j = 1 \dots 13$,
 $d_{8,j,2k-1} = \rho(h_{13A})_{\nu^j}(h_{7A} \cdot \vartheta_k)$ for $k = 1, 2$ and $j = 1 \dots 13$,
 $d_{8,j,2k} = \rho(h_{13A})_{\nu^j}(h_{7A}^2 \cdot \vartheta_k)$ for $k = 1, 2$ and $j = 13$.

Acknowledgments. Some calculations were carried out at the KanBalam supercomputer of DGTIC–UNAM.

References

- [1] Arbarello, E.; Cornalba, M.; Griffiths, P. A.; Harris, J. *Geometry of algebraic curves*. Vol. I. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 267. Springer-Verlag, New York, 1985.
- [2] Conder, Marston *The genus of compact Riemann surfaces with maximal automorphism group*. J. Algebra 108 (1987), no. 1, 204–247.

- [3] Cooperman, Gene. *Parallel GAP/MPI (ParGAP/MPI)*, Version 1, College of Computer Science, Northeastern University, 1999, <http://www.ccs.neu.edu/home/gene/pargap.html>.
- [4] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4.12*; 2008, <http://www.gap-system.org>.
- [5] Harris, Joe; Morrison, Ian. *Moduli of curves*. Graduate Texts in Mathematics, 187. Springer-Verlag, New York, 1998.
- [6] A. M. Macbeath, *Action of automorphisms of a compact Riemann surface on the first homology group*, Bull. London Math. Soc.5 (1973) 103–108.
- [7] Moreno-Mejía, Israel The quadrics through the Hurwitz curves of genus 14. J. Lond. Math. Soc. (2) 81 (2010), no. 2, 374–388.
- [8] Mumford. D. *Curves and their Jacobians*. The University of Michigan Press, Ann Arbor, Mich., 1975.
- [9] Petri. K, *Über die invariante Darstellung algebraischer Funktionen einer Veränderlichen*. Math. Ann. 88, 242 (1922).
- [10] Saint-Donat, B. *On Petri’s analysis of the linear system of quadrics through a canonical curve*. Math. Ann. 206 (1973), 157–175
- [11] Sinkov, Abraham On the group-defining relations $(2, 3, 7; p)$. Ann. of Math. (2) 38 (1937), no. 3, 577–584.
- [12] Streit, Manfred *Field of definition and Galois orbits for the Macbeath-Hurwitz curves*. Arch. Math. 74 (2000), no. 5, 342–49.
- [13] Völklein, Helmut. *Groups as Galois groups*. An introduction. Cambridge Studies in Advanced Mathematics, 53. Cambridge University Press, Cambridge, 1996.

Israel Moreno Mejía.
 Instituto de Matemáticas
 Universidad Nacional Autónoma de México
 México, D.F. C.P. 04510
 México.

E-mail address: **israel@math.unam.mx**